



Lesson 8: Similarity

Student Outcomes

- Students know the definition of similar and why dilation alone is not enough to determine similarity.
- Given two similar figures, students describe the sequence of a dilation and a congruence that would map one figure onto the other.

Lesson Notes


In Module 2, students used vectors to describe the translation of the plane. Now in Topic B, figures are bound to the coordinate plane, and students will describe translations in terms of units left or right and/or units up or down. When figures on the coordinate plane are rotated, the center of rotation is the origin of the graph. In most cases, students will describe the rotation as having center O and degree d unless the rotation can be easily identified, i.e., a rotation of 90° or 180° . Reflections remain reflections across a line, but when possible, students should identify the line of reflection as the x -axis or y -axis.

It should be noted that congruence, together with similarity, is *the* fundamental concept in planar geometry. It is a concept defined without coordinates. In fact, it is most transparently understood when introduced without the extra conceptual baggage of a coordinate system. This is partly because a coordinate system picks out a preferred point (the origin), which then centers most discussions of rotations, reflections, and translations at or in reference to that point. They are then further restricted to only the "nice" rotations/reflections/translations that are easy to do in a coordinate plane. Restricting to "nice" transformations is a huge mistake mathematically because it is antithetical to the main point that must be made about congruence: that rotations, translations, and reflections are abundant in the plane—that for every point in the plane, there are an *infinite number* of rotations up to 360° , that for every line in the plane there is a reflection, and for every directed line segment there is a translation. It is this abundance that helps students realize that every congruence transformation (i.e., the act of "picking up a figure" and moving it to another location) can be accomplished through a sequence of translations, rotations, and reflections, and further, that similarity is a congruence transformation in addition to dilation.

Classwork

Concept Development (5 minutes)

- A dilation alone is not enough to state that two figures are similar. Consider the following pair of figures:



- Do these figures look similar?
 - *Yes, they look like the same shape, but they are different in size.*

- How could you prove that they are similar? What would you need to do?
 - *We would need to show that they could become the same size by dilating one of the figures.*
- Would we be able to dilate one figure so that it was the same size as the other?
 - *Yes, we could dilate to make them the same size by using the appropriate scale factor.*
- We could make them the same size, but would a dilation alone map figure S onto S_0 ?
 - *No, a dilation alone would not map S onto S_0 .*
- What else should we do to map figure S onto S_0 ?
 - *We would have to perform a translation and a rotation to map S onto S_0 .*
- That is precisely why a dilation alone is not enough to define similarity. Two figures are said to be similar if one can be mapped onto the other using a dilation followed by a congruence (a sequence of basic rigid motions) or a congruence followed by a dilation.

Example 1 (5 minutes)

Example 1


In the picture below we have a triangle ABC , that has been dilated from center O , by a scale factor of $r = \frac{1}{2}$. It is noted by $A'B'C'$. We also have triangle $A''B''C''$, which is congruent to triangle $A'B'C'$ (i.e., $\triangle A'B'C' \cong \triangle A''B''C''$).

Describe the sequence that would map triangle $A''B''C''$ onto triangle ABC .

Scaffolding:
Remind students of the work they did in Lesson 3 to bring dilated figures back to their original size.

- Based on the definition, how could we show that triangle $A''B''C''$ is similar to triangle ABC ?
 - *To show that $\triangle A''B''C'' \sim \triangle ABC$, we need to describe a dilation followed by a congruence.*

- We want to describe a sequence that would map triangle $A''B''C''$ onto triangle ABC . There is no clear way to do this so let's begin with something simpler: How can we map triangle $A'B'C'$ onto triangle ABC ? That is, what is the precise dilation that would make triangle $A'B'C'$ the same size as triangle ABC ?
 - A dilation from center O with scale factor $r = 2$.
- Remember, our goal was to describe how to map triangle $A''B''C''$ onto triangle ABC . What precise dilation would make triangle $A''B''C''$ the same size as triangle ABC ?
 - A dilation from center O with scale factor $r = 2$ would make triangle $A''B''C''$ the same size as triangle ABC .
- (Show the picture below with the dilated triangle $A''B''C''$ noted by $A'''B'''C'''$.) Now that we know how to make triangle $A''B''C''$ the same size as triangle ABC , what rigid motion(s) should we use to actually map triangle $A''B''C''$ onto triangle ABC ? Have we done anything like this before?




- Number 2 of the problem set from Lesson 2 was like this. That is, we had two figures dilated by the same scale factor in different locations on the plane. To get one to map to the other, we just translated along a vector.
- Now that we have an idea of what needs to be done, let's describe the translation in terms of coordinates. How many units and in which direction will we need to translate so that triangle $A'''B'''C'''$ maps to triangle ABC ?
 - We need to translate triangle $A'''B'''C'''$ 20 units to the left, and 2 units down.
- Let's use precise language to describe how to map triangle $A''B''C''$ onto triangle ABC . We will need information about the dilation and the translation.
 - The sequence that would map triangle $A''B''C''$ onto triangle ABC is as follows: Dilate triangle $A''B''C''$ from center O by scale factor $r = 2$. Then translate the dilated triangle 20 units to the left and 2 units down.
- Since we were able to map triangle $A''B''C''$ onto triangle ABC with a dilation followed by a congruence, we can write that triangle $A''B''C''$ is similar to triangle ABC , in notation, $\Delta A''B''C'' \sim \Delta ABC$.

MP.6

Example 2 (5 minutes)

- In the picture below, we have a triangle DEF , that has been dilated from center O , by scale factor $r = 3$. It is noted by $D'E'F'$. We also have a triangle $D''E''F''$, which is congruent to triangle $D'E'F'$ (i.e., $\triangle D'E'F' \cong \triangle D''E''F''$).



- We want to describe a sequence that would map triangle $D''E''F''$ onto triangle DEF . This is similar to what we did in the last example. Can someone summarize the work we did in the last example?


 - First, we figured out what scale factor r would make the triangles the same size. Then, we used a translation to map the magnified figure onto the original triangle.
- What is the difference between this problem and the last?

 - This time the scale factor is greater than one, so we will need to shrink triangle $D''E''F''$ to the size of triangle DEF . Also, it appears as if a translation alone will not map one triangle onto another.
- Now, since we want to dilate triangle $D''E''F''$ to the size of triangle DEF , we need to know what scale factor r is required so that $|OF| = r|OF''|$. What scale factor do you think we should use and why?

 - We need a scale factor $r = \frac{1}{3}$ because we want $|OF| = r|OF''|$.
- What precise dilation would make triangle $D''E''F''$ the same size as triangle DEF ?


 - A dilation from center O with scale factor $r = \frac{1}{3}$ would make triangle $D''E''F''$ the same size as triangle DEF .

- (Show the picture below with the dilated triangle $D''E''F''$ noted by $D'''E'''F'''$.) Now, we should use what we know about rigid motions to map the dilated version of triangle $D''E''F''$ onto triangle DEF . What should we do first?



▫ We should translate triangle $D'''E'''F'''$ 2 units to the right.

- (Show the picture below, the translated triangle is noted in red.) What should we do next (refer to the translated triangle as the red triangle)?




▫ Next, we should reflect the red triangle across the x -axis to map the red triangle onto triangle DEF .

MP.6


- Use precise language to describe how to map triangle $D''E''F''$ onto triangle DEF .
 - *The sequence that would map triangle $D''E''F''$ onto triangle DEF is as follows: Dilate triangle $D''E''F''$ from center O by scale factor $r = \frac{1}{3}$. Then translate the dilated image of triangle $D''E''F''$, noted by $D'''E'''F'''$ two units to the right. Finally, reflect across the x -axis to map the red triangle onto triangle DEF .*
- Since we were able to map triangle $D''E''F''$ onto triangle DEF with a dilation followed by a congruence, we can write that triangle $D''E''F''$ is similar to triangle DEF . (In notation: $\triangle D''E''F'' \sim \triangle DEF$.)

Example 3 (3 minutes)

- In the diagram below, $\triangle ABC \sim \triangle A'B'C'$. Describe the sequence of the dilation followed by a congruence that would prove these figures to be similar.




- Let's begin with the scale factor. We know that $r|AB| = |A'B'|$. What scale factor r will make $\triangle ABC$ the same size as $\triangle A'B'C'$?
 - *We know that $r \times 2 = 1$; therefore, $r = \frac{1}{2}$ will make $\triangle ABC$ the same size as $\triangle A'B'C'$.*
- If we apply a dilation from the origin of scale factor $r = \frac{1}{2}$, then the triangles will be the same size (as shown and noted by triangle $A''B''C''$). What sequence of rigid motions would map the dilated image of $\triangle ABC$ onto $\triangle A'B'C'$?




- We could translate the dilated image of $\triangle ABC$, $\triangle A''B''C''$, 3 units to the right and 4 units down, then reflect the triangle across line $A'B'$.
- The sequence that would map $\triangle ABC$ onto $\triangle A'B'C'$ to prove the figures similar is: A dilation from the origin by scale factor $r = \frac{1}{2}$, followed by the translation of the dilated version of $\triangle ABC$ 3 units to the right and 4 units down, followed by the reflection across line $A'B'$.

Example 4 (5 minutes)

- In the diagram below, we have two similar figures. Using the notation, we have $\triangle ABC \sim \triangle DEF$. We want to describe the sequence of the dilation followed by a congruence that would prove these figures to be similar.



- First, we need to describe the dilation that would make the triangles the same size. What information do we have to help us describe the dilation?
 - *Since we know the length of side AC and side DF, we can determine the scale factor.*
- Can we use any two sides to calculate the scale factor? Assume, for instance, that we know that side AC is 18 units in length and side EF is 2 units in length. Could we find the scale factor using those two sides, AC and EF? Why or why not?
 - *No. We need more information about corresponding sides. Sides AC and DF are the longest sides of each triangle (they are also opposite the obtuse angle in the triangle). Side AC does not correspond to side EF. If we knew the length of BC, we could use BC and EF.*
- Now that we know that we can find the scale factor if we have information about corresponding sides, how would we calculate the scale factor if we were mapping $\triangle ABC$ onto $\triangle DEF$?
 - $|DF| = r|AC|$, then $6 = r \times 18$, and $r = \frac{1}{3}$.
- If we were mapping $\triangle DEF$ onto $\triangle ABC$, what would the scale factor be?
 - $|AC| = r|DF|$, then $18 = r \times 6$, and $r = 3$.
- What is the precise dilation that would map $\triangle ABC$ onto $\triangle DEF$?
 - *Dilate $\triangle ABC$ from center O , by scale factor $r = \frac{1}{3}$.*
- (Show the picture below with the dilated triangle noted as $\triangle A'B'C'$.) Now we have to describe the congruence. Work with a partner to determine the sequence of rigid motions that would map $\triangle ABC$ onto $\triangle DEF$.




- *Translate the dilated version of $\triangle ABC$ 7 units to the right and 2 units down. Then, rotate d degrees around point E , so that segment $B'C'$ maps onto segment EF . Finally, reflect across line EF .*

Note that “ d degrees” refers to a rotation by an appropriate number of degrees to exhibit similarity. Students may choose to describe this number of degrees in other ways.

- The sequence of a dilation followed by a congruence that proves $\triangle ABC \sim \triangle DEF$ is as follows: Dilate $\triangle ABC$ from center O by scale factor $r = \frac{1}{3}$. Translate the dilated version of $\triangle ABC$ 7 units to the right and 2 units down. Next, rotate around point E d degrees so that segment $B'C'$ maps onto segment EF , then reflect the triangle across line EF .

Example 5 (3 minutes)


- Knowing that a sequence of a dilation followed by a congruence defines similarity also helps determine if two figures are, in fact, similar. For example, would a dilation map triangle ABC onto triangle DEF ? That is, is $\triangle ABC \sim \triangle DEF$?



- No, by FTS, we expect the corresponding sides lengths to be in proportion and equal to the scale factor. If we compare side AC to side DF , and BC to EF , then we get $\frac{18}{6} \neq \frac{15}{4}$.
- Therefore, the triangles are not similar, because a dilation will not map one to the other.

Example 6 (3 minutes)

- Again, knowing that a dilation followed by a congruence defines similarity also helps determine if two figures are, in fact, similar. For example, would a dilation map Figure A onto Figure A' ? That is, is $Figure A \sim Figure A'$?




- No, even though we could say that the corresponding sides are in proportion, there exists no single rigid motion or sequence of rigid motions that would map a four-sided figure to a three-sided figure. Therefore, the figures do not fulfill the congruence part of the definition for similarity, and Figure A is not similar to Figure A' .

Exercises 1–4 (10 minutes)

Allow students to work in pairs to describe sequences that map one figure onto another.


Exercises 1–4

- Triangle ABC was dilated from center O by scale factor $r = \frac{1}{2}$. The dilated triangle is noted by $A'B'C'$. Another triangle $A''B''C''$ is congruent to triangle $A'B'C'$ (i.e., $\triangle A''B''C'' \cong \triangle A'B'C'$). Describe the dilation followed by the basic rigid motion that would map triangle $A''B''C''$ onto triangle ABC .




Triangle $A''B''C''$ will need to be dilated from center O , by scale factor $r = 2$ to bring it to the same size as triangle ABC . This will produce a triangle noted by $A'''B'''C'''$. Next, triangle $A'''B'''C'''$ will need to be translated 4 units up and 12 units left. The dilation followed by the translation will map triangle $A''B''C''$ onto triangle ABC .

2. Describe the sequence that would show $\triangle ABC \sim \triangle A'B'C'$.



Since $r|AB| = |A'B'|$, then $r \times 2 = 6$ and $r = 3$. A dilation from the origin by scale factor $r = 3$ will make $\triangle ABC$ the same size as $\triangle A'B'C'$. Then, a translation of the dilated image of $\triangle ABC$ ten units right and five units down, followed by a rotation of 90 degrees around point C' will map $\triangle ABC$ onto $\triangle A'B'C'$, proving the triangles to be similar.

3. Are the two triangles shown below similar? If so, describe the sequence that would prove $\triangle ABC \sim \triangle A'B'C'$. If not, state how you know they are not similar.




Yes, triangle $\triangle ABC \sim \triangle A'B'C'$. The corresponding sides are in proportion, and equal to the scale factor:

$$\frac{10}{15} = \frac{4}{6} = \frac{12}{18} = \frac{2}{3} = r.$$

To map triangle ABC onto triangle $A'B'C'$, dilate triangle ABC from center O , by scale factor $r = \frac{2}{3}$. Then, translate triangle ABC along vector $\overrightarrow{AA'}$. Next, rotate triangle ABC d degrees around point A .

4. Are the two triangles shown below similar? If so, describe the sequence that would prove $\triangle ABC \sim \triangle A'B'C'$. If not, state how you know they are not similar.



Yes, triangle $\triangle ABC \sim \triangle A'B'C'$. The corresponding sides are in proportion and equal to the scale factor:

$$\frac{4}{3} = \frac{8}{6} = \frac{4}{3} = 1.3\bar{3}; \frac{10.67}{8} = 1.33375; \text{ therefore, } r = 1.33 \cong \frac{4}{3}$$

To map triangle ABC onto triangle $A'B'C'$, dilate triangle ABC from center O , by scale factor $r = \frac{4}{3}$. Then, translate triangle ABC along vector $\overrightarrow{AA'}$. Next, rotate triangle ABC 180 degrees around point A' .

Closing (5 minutes)

Summarize, or ask students to summarize, the main points from the lesson:

- We know that similarity is defined as the sequence of a dilation, followed by a congruence.
- To show that a figure in the plane is similar to another figure of a different size, we must describe the sequence of a dilation, followed by a congruence (one or more rigid motions), that maps one figure onto another.

Lesson Summary

Similarity is defined as mapping one figure onto another as a sequence of a dilation followed by a congruence (a sequence of rigid motions).

The notation, $\triangle ABC \sim \triangle A'B'C'$, means that $\triangle ABC$ is similar to $\triangle A'B'C'$.

Exit Ticket (5 minutes)


Name _____

Date _____

Lesson 8: Similarity


Exit Ticket

In the picture below, we have a triangle DEF that has been dilated from center O , by scale factor $r = \frac{1}{2}$. The dilated triangle is noted by $D'E'F'$. We also have a triangle $D''EF$, which is congruent to triangle DEF (i.e., $\triangle DEF \cong \triangle D''EF$). Describe the sequence of a dilation followed by a congruence (of one or more rigid motions) that would map triangle $D'E'F'$ onto triangle $D''EF$.



Exit Ticket Sample Solutions

In the picture below we have a triangle DEF that has been dilated from center O , by scale factor $r = \frac{1}{2}$. The dilated triangle is noted by $D'E'F'$. We also have a triangle $D''E'F'$, which is congruent to triangle DEF (i.e., $\triangle DEF \cong \triangle D''E'F'$). Describe the sequence of a dilation, followed by a congruence (of one or more rigid motions), that would map triangle $D'E'F'$ onto triangle $D''E'F'$.




Triangle $D'E'F'$ will need to be dilated from center O , by scale factor $r = 2$ to bring it to the same size as triangle DEF . This will produce the triangle noted by DEF . Next, triangle DEF will need to be reflected across line EF . The dilation followed by the reflection will map triangle $D'E'F'$ onto triangle $D''E'F'$.

Problem Set Sample Solutions


Students practice dilating a curved figure and describing a sequence of a dilation followed by a congruence that maps one figure onto another.

- In the picture below, we have a triangle DEF , that has been dilated from center O , by scale factor $r = 4$. It is noted by $D'E'F'$. We also have a triangle $D''E''F''$, which is congruent to triangle $D'E'F'$ (i.e., $\triangle D'E'F' \cong \triangle D''E''F''$). Describe the sequence of a dilation, followed by a congruence (of one or more rigid motions), that would map triangle $D''E''F''$ onto triangle DEF .




First, we must dilate triangle $D''E''F''$ by scale factor $r = \frac{1}{4}$ to shrink it to the size of triangle DEF . Next, we must translate the dilated triangle, noted by $D'''E'''F'''$, one unit up and two units to the right. This sequence of the dilation followed by the translation, would map triangle $D''E''F''$ onto triangle DEF .

2. Triangle ABC was dilated from center O by scale factor $r = \frac{1}{2}$. The dilated triangle is noted by $A'B'C'$. Another triangle $A''B''C''$ is congruent to triangle $A'B'C'$ (i.e., $\Delta A''B''C'' \cong \Delta A'B'C'$). Describe the dilation followed by the basic rigid motion that would map triangle $A''B''C''$ onto triangle ABC .




Triangle $A''B''C''$ will need to be dilated from center O by scale factor $r = 2$ to bring it to the same size as triangle ABC . This will produce a triangle noted by $A'''B'''C'''$. Next, triangle $A'''B'''C'''$ will need to be translated 18 units to the right and 2 units down, producing the triangle shown in red. Next, rotate the red triangle d degrees around point B , so that one of the segments of the red triangle coincides completely with segment BC . Then, reflect the red triangle across line BC . The dilation, followed by the congruence described, will map triangle $A''B''C''$ onto triangle ABC .

3. Are the two figures shown below similar? If so, describe the sequence that would prove the similarity. If not, state how you know they are not similar.



No, these figures are not similar. There is no single rigid motion, or sequence of rigid motions, that would map Figure A onto Figure B.


4. Triangle ABC is similar to triangle $A'B'C'$, (i.e., $\Delta ABC \sim \Delta A'B'C'$). Prove the similarity by describing the sequence that would map triangle A



The scale factor that would prove the similarity of the triangles is a dilation from center O by a scale factor of $r = 3$, followed by a translation along vector $\overrightarrow{A'A}$ and, finally, a reflection across line AC .


3. The sequence that would prove the similarity of the triangles is a dilation from center O by a scale factor of $r = 3$, followed by a translation along vector $\overrightarrow{A'A}$ and, finally, a reflection across line AC .

5. Are the two figures shown below similar? If so, describe the sequence that would prove $\triangle ABC \sim \triangle A'B'C'$. If not, state how you know they are not similar.



Yes, the triangles are similar. The scale factor that triangle ABC has been dilated is $r = \frac{1}{5}$. The sequence that proves the triangles are similar is as follows: dilate triangle $A'B'C'$ from center O by scale factor $r = 5$, then translate triangle $A'B'C'$ along vector $\vec{C'C}$, next, rotate triangle $A'B'C'$ d degrees around point C and, finally, reflect triangle $A'B'C'$ across line AC .

6. Describe the sequence that would show $\triangle ABC \sim \triangle A'B'C'$.



Since $r|AB| = |A'B'|$, then $r \times 3 = 1$, and $r = \frac{1}{3}$. A dilation from the origin by scale factor $r = \frac{1}{3}$ will make $\triangle ABC$ the same size as $\triangle A'B'C'$. Then, a translation of the dilated image of $\triangle ABC$ four units down and one unit to the right, followed by a reflection across line $A'B'$ will map $\triangle ABC$ onto $\triangle A'B'C'$, proving the triangles to be similar.